

K. with preface

THEOREMATUM
IN LIBRIS
ARCHIMEDIS
DE
SPHAERA & CYLINDRO
Declaratio.

Authore
GUILIELMO OUGHTREDO
ANGL.

OXONIAE,
Excudebat LEON. LICHFIELD, Veneunt
apud THO. ROBINSON. Anno
Dom. 1652.

THE ORE MOUNTAIN

IN THE

ARCHIVES

OF THE

DEPARTMENT

OF THE

OF THE

OF THE

OF THE

OF THE



Rerum quarundam denotationes.

R radius, est semidiameter circuli, siue uno constet nomine AO, vel $E\omega$, vel IU : siue duobus ut AO + $E\omega$, vel $E\omega$ + IU : ut in schemate 1.

Δ . π :: semidiameter. semiperipheria.

$\frac{\pi}{\Delta}$ R, est semiperipheria circuli cuius Radius est R.

$\frac{\pi}{\Delta}$: AO + $E\omega$: est semiperipheria circuli cuius Radius est AO + $E\omega$.

$\frac{\pi}{\Delta}$ Rq, est area circuli.

$\frac{\pi}{3\Delta}$ Rq * altitud: vel $\frac{\pi}{\Delta}$ Rq in $\frac{1}{3}$ Altitud, est Conus; scil: $\frac{1}{3}$ Cylindri.

○ significat superficiem curvam.

Coni & Cylindri, qui in æqualibus sunt basibus, sunt ut altitudines. 14 e 12.

Æqualium Conorum & Cylindrorum bases & altitudines reciprocantur. 15 e 12.

Assumo, Figuram regularem infinitorum laterum, cui nec major inscribi, nec minor circumscribi poterit; si plana sit, esse circulum; sin solida, esse sphaeram.

Theorematum



Theorematum in Libris ARCHIMEDIS de Sphæra & Cylindro

DECLARATIO.

DUODECIM primas propositiones, quia demonstrationibus negativis, quas ego ut parum scientificas, quantum possum, evito, inque ipsarum loco affirmativas substituo, inserviunt, missas faciam.

I. In Cylindro recto, Si $2R, M,$ Latus; hoc est, $2AO, M, KA,$:: Dico $\frac{\pi}{\int} Mq = \bigcirc$ Cylindri.

Nam $\frac{\pi}{\int} Mq = \frac{\pi}{\int} 2AO \times KA.$ 13 l 1.

(Ad septem theoremata sequentia pertinet schema I.)

II. In Cono æquicruro KON, si KO, M, AO :: Dico $\frac{\pi}{\int} Mq = \bigcirc$ Coni. Nam $\frac{\pi}{\int} Mq = \frac{\pi}{\int} AO$ in KO. 14 l 1.

A 3

III In

2 De Sphæra & Cylindro.

III. In Cono æquicruro KON, Dico esse semid:
balis. Latus :: Basis. O Coni. Nam AO. KO ::
 $\frac{\pi}{\delta}$ AOq. $\frac{\pi}{\delta}$ AO in KO. 15 l 1.

IV. In Cono æquicruro KON, Si AO+E ω , M,
O ω =KO-K ω :: Dico O frusti O ω N=($\frac{\pi}{\delta}$ Mq) $\frac{\pi}{\delta}$:
AO+E ω : *O ω . Nam per 2, O O ω N= $\frac{\pi}{\delta}$: AO*KO:
mi $\frac{\pi}{\delta}$: E ω *K ω = $\frac{\pi}{\delta}$: AO+E ω : in: KO-K ω . Est enim
AO+E ω in KO-K ω =AO * KO -E ω *K ω pl E ω *KO
-AO*KO, quæ se invicem tollunt: Quia AO.E ω ::
KO.K ω .

V. In Cono æquicruro KON, Si KO, M, AO ::;
& AP perpendicularis lateri KO: Dico ($\frac{\pi}{3\delta}$ Mq)
 $\frac{\pi}{3\delta}$ AO*KO in AP= $\frac{\pi}{3\delta}$ AOq in KA=KON. Nam
KA. AP:: KO. AO:: AO*KO. AOq. Ergo.
17 l 1.

VI. In Cono æquicruro KON, Si K ω . M. E ω ::;
& AP perpend: lateri KO: Dico ($\frac{\pi}{3\delta}$ Mq) $\frac{\pi}{3\delta}$ E ω
*K ω in AP= $\frac{\pi}{3\delta}$ E ω q*K ω , scil: rhombo K ω A ν . Nam
KA. AP:: K ω . E ω :: E ω *K ω . E ω q. Ergo.
18 l 1.

VII. In Cono æquicruro KON, Dico frustum
Conicè

Conicè excavatum $O\omega A\nu N$, æquari Cono cujus basis est æqualis \bigcirc frusti $O\omega\nu N$, & altitudo AP : hoc est,

con: KON — rhomb: $K\omega A\nu = \frac{\pi}{\delta}$: $AO + E\omega$: * $O\omega$ in

$\frac{1}{2}AP$. Nam $\frac{\pi}{\delta} AO * KO$ in $\frac{1}{2}AP = KON$, per 5

Et $\frac{\pi}{\delta} E\omega * K\omega$ in $\frac{1}{2}AP = \text{rhomb: } K\omega A\nu$, per 6 } ho-

rum differentia est $\frac{\pi}{\delta} AO * KO - \frac{\pi}{\delta} E\omega * K\omega$, per 4,

$= \frac{\pi}{\delta}$: $AO + E\omega$: * $O\omega = \bigcirc O\omega A\nu$; ductis omnibus in

$\frac{1}{2}AP$. Ergo, &c.

VIII. In Cono æquicruro KON , Dico rhombum conicè excavatum $\omega UAS\nu$, æquari Cono cujus basis est æqualis \bigcirc frusti $\omega U S\nu$, & altitudo AP : hoc est, rhomb $K\omega A\nu$ — rhomb


$KUAS = \frac{\pi}{\delta}$: $E\omega + IU$:

* ωU in $\frac{1}{2}AP$. Nam

per 6, $\frac{\pi}{\delta} E\omega * K\omega$ in $\frac{1}{2}AP = \text{rhomb } K\omega A\nu$ } horum

Et $\frac{\pi}{\delta} IU * KU$ in $\frac{1}{2}AP = \text{rhomb } KUAS$ }

differentia est $\frac{\pi}{\delta} E\omega * K\omega - \frac{\pi}{\delta} IU * KU$, per 4, $= \frac{\pi}{\delta}$: $E\omega$



4


De Sphæra & Cylindro.

$E\omega + IU: \kappa\omega U = O\omega US$; ductis omnibus in $\frac{1}{2}AP$.
Ergo, &c.

IX. Si figura plana polygona laterum æqualium
& numero parium. $ABDF\delta\beta\alpha\gamma\epsilon\phi EC$, inscribatur cir-
culo, junganturque anguli rectis lineis parallelis:
Dico $AB.B\alpha :: A\alpha.BC + DE + F\phi + \delta\epsilon + \beta\gamma$; hoc est, $2BC$
 $+ 2DE + F\phi$. Nam $AB.B\alpha :: \frac{1}{2}AK.$ $\frac{1}{2}BC :: \frac{1}{2}KL.$ $\frac{1}{2}DE ::$
 $\frac{1}{2}L\lambda.$ $\frac{1}{2}F\phi :: \frac{1}{2}\lambda\kappa.$ $\frac{1}{2}\delta\epsilon :: \frac{1}{2}\kappa\alpha.$ $\frac{1}{2}\beta\gamma$.

Quare $A\alpha \times B\alpha = AB$ in $2BC + 2DE + L\phi$. 2111.

Et in segmento $A\delta\epsilon$, erit $AB.B\alpha :: A\eta.BC + DE$
 $+ F\phi + \frac{1}{2}\delta\epsilon$.



Quare

De Sphæra & Cylindro.

5

Quare $A\alpha \times B\alpha = AB$ in $BC + DE + F\phi + \frac{1}{2} \delta \epsilon$.

X. Si circulo, vel circuli segmento alicui figura ejusmodi plana polygona laterum æqualium & parium, tum inscribatur, tum circumscribatur; & diametro $A\alpha$ quiescente, circulus circumvolvatur; describetur figura solida constans superficiebus quibusdam Conicis: Et paralleli $BC, DE, F\phi, \delta \epsilon, \beta \gamma$, describent totidem circulos parallelos. Atque in his, quæ circumscripta est, sive continens, major semper est circulo incluso: & quæ inscripta est, minor semper erit circulo ambiente. Et superficies figuræ circumscriptæ, ad superficiem figuræ inscriptæ similis, est in ratione laterum duplicata: At figura ipsa solida circumscripta, ad solidam similem inscriptam, in ratione triplicata. 22. 27. 30. 34. 37 l 1.

XI. Si diameter circuli includentis ejusmodi figuram solidam, sit $A\alpha$: fiatque $A\alpha, M, B\alpha ::$ vel, quod idem est, per 9, $2BC + 2DE + F\phi, M, AB ::$ Dico

$\frac{\pi}{\delta} Mq =$ superficiei figuræ. Nam per 2, $\frac{\pi}{\delta} BC \times AB$

$= 2 \circ$ coni ABC : & per 4, $\frac{\pi}{\delta} BC + DE$ in $AB = 2 \circ$

frusti $BCED$: & $\frac{\pi}{\delta} DE + F\phi$ in $AB = 2 \circ$ frusti $DE\phi F$.

Ergo $\frac{\pi}{\delta} 2BC + 2DE + F\phi$ in $AB = \circ$ figuræ totius,

nempe $\frac{\pi}{\delta} Mq$. 23. 28 l 1.

XII. In Schem: 3. Figuræ ejusmodi solidæ, si
sphæræ

sphæræ inscribatur, superficies $\frac{\pi}{3} Mq$ minor est circulo habente axem sphæræ continentis Aa pro diametro. Nam $M \sqsubset Aa$.

Sin circumscribatur, superficies $\frac{\pi}{3} Mq$ major est circulo habente axem sphæræ contentæ $2IP = Ba$ pro diametro. Nam $Aa, M, 2IP \div \div$: Quare M cadet inter A & Q . 24, 29 l 1.

XIII. Quidni igitur sphæræ superficies æquetur quatuor maximis circulis; nempe $\frac{\pi}{3} Diam : q?$

31 l 1.

XIV. Figura ejusmodi solida æqualis est Cono, cujus Basis est circulus æqualis superficiei figuræ; & Altitudo IP perpenditè centro sphæræ in latus figuræ: hoc est, per 11, $\frac{\pi}{3} Mq$ in $(IP) \frac{1}{2} Ba =$ figuræ toti solidæ. Nam per 6, Rhomb: $BACI = \frac{1}{3} \cap BAC$ in IP . Et per 8, Excavatum $DBCE = \frac{1}{3} \cap DBCE$ in IP . Et per 7, Excavatum $FDIE = \frac{1}{3} \cap FDE$ in IP . Et similiter pro altero hæmisphærio. Quare $\frac{1}{3} \cap BAC + \frac{1}{3} \cap DBCE + \frac{1}{3} \cap FDE$ in $Ba (2IP) =$ toti figuræ solidæ; nempe $\frac{\pi}{3} Mq$: vel $\frac{\pi}{3} Aa \times Ba$ in $\frac{1}{2} BA(IP)$.

25, 29 l 1.

XV. Figura ejusmodi, si sphæræ inscribatur, minor est quatuor Conis habentibus basem æqualem circulo sphæræ maximo: hoc est Cono habenti basem æqualem superficiei sphæræ; altitudinem verò æqualem semiaxi. Sin circumscribatur, iisdem major est.

Nam

De Sphæra & Cylindro. 7

Nam per 12, superficies figuræ inscriptæ, superficie sphæræ minor est: circumscriptæ autem, major. 26.
29 l 1.

XVI. Quidni igitur ipsa sphæra æqualis sit quatuor Conis habentibus basem æqualem circulo sphæræ maximo; hoc est Cono habenti basem æqualem superficiei sphæræ; Altitudinem verò æqualem semi-axi? 32 l 1.

Conject. $\frac{2}{3}$ Cylind: = Sphæræ = 2 Conis. Nam

$$\frac{\pi}{3\delta} Rq + R = \text{Sphæræ. Et } \frac{\pi}{\delta} Rq \times 2R = \text{Cylindro.}$$

$$\frac{\pi}{3\delta} Rq \times 2R = \text{Cono.}$$

XVII. Si figura ejusmodi sive inscribatur, sive circumscribatur, segmento sphæræ, puta $A\delta\epsilon$, cujus balis sit $\delta\epsilon$; altitudo $A\eta$; fiatque $B\alpha$, M , $A\eta$:: vel, quod idem est, per 9, $BC + DE + F\phi + \frac{1}{2}\delta\epsilon$, M , AB :: Dico

$$\frac{\pi}{\delta} Mq = \text{superficiei figuræ illius mancæ. Nam per 2,}$$

$$\frac{\pi}{\delta} \frac{1}{2}BC \text{ in } AB = \bigcirc ABC. \text{ Et per 4, } \frac{\pi}{\delta} \frac{1}{2}BC + \frac{1}{4}DE$$

$$\text{in } AB = \bigcirc DBCE: \text{ Et } \frac{\pi}{\delta} \frac{1}{2}DE + \frac{1}{2}F\phi \text{ in } AB = \bigcirc FDE\phi.$$

$$\text{Et } \frac{\pi}{\delta} \frac{1}{2}F\phi + \frac{1}{2}\delta\epsilon \text{ in } AB = \bigcirc \delta F\phi\epsilon. \text{ Ergo 33. 37 l 1.}$$


XVIII. Figuræ ejusmodi mancæ, si segmento sphæræ, puta $A\delta\epsilon$, inscribatur, superficies $\frac{\pi}{\delta} Mq$

$$\sqsubset \frac{\pi}{\delta} A\delta q. \text{ Nam } A\delta q = A\alpha \times A\eta \sqsubset E\alpha \times A\eta.$$

Sin

Sin circumscribatur, superficies $\frac{\pi}{2} Mq = \frac{\pi}{2} Aq$.

Nam $Ba = 2IQ$ est diameter sphære interioris sive contentæ. Estque $An = Qn$. Quare $\frac{1}{2} M$ protenditur ultra IQ diametrum sphære contentæ. 35.38.41 l 1.



XIX. Quidni igitur ipsa superficies segmenti sphære æqualis sit circulo, cuius semidiameter est recta ducta a vertice segmenti in finem basis? 40 l 1.

XX. Figura ejusmodi manca, sive inscribatur segmento sphære, puta DAE minori semicirculo, vel DaE majori, æqualis est Cono habenti basem æqualem superfici ei illius figuræ mancæ; altitudinem
verò

De Sphæra & Cylindro.

9

verò æqualem perpend : IP è centro in latus figuræ ,
 adjecto vel ablato Cono DIE medio ; hoc est , toti
 solido DBACEI , vel reliquo DFδβαγεφEI. Nam in
 solido minore DBA=CEI , per 6 , Rhomb: BACI
 = $\frac{1}{3}$ ∩ BAC in IP. Et per 8 , $\frac{1}{3}$ ∩ DBCE in IP =
 excavato DBICE. Ergo. Et similiter de majore
 DFδβαγεφEI. 36. 39 l i.

Consect. Quare & per 18 , Si figura ejusmodi seg-
 mento sphæræ, puta DAE, vel DαE, inscribatur, totum
 solidum minus est Cono, cujus basis est circulus a se-
 midiametro AD, vel αD ; & altitudo semiaxis IA.

XXI. Quidni igitur Sector sphæræ DBACEI , æ-
 qualis sit Cono $\frac{\pi}{3\delta}$ ADq in IA ; & reliquus Dα=
 EI, æqualis Cono $\frac{\pi}{3\delta}$ αDq in IA? 42 l i.

XXII. Si fiat Hα. Hα+Iα.:

HA. HS: Dico $\frac{\pi}{3\delta}$ DHq in

HS=segm: sphæræ DAE.

Nam Hα. HA:: Hα + Iα

-Hα. HS-HA:: Iα. AS.

Estque (Iα+AS) IS. IA::


(Hα+HA) Aα. Hα:: Aαq.

αDq:: ADq. DHq. Quare

$\frac{\pi}{3\delta}$ DHq * IS = $\frac{\pi}{3\delta}$ ADq

* IA = segmento DAE†

(Cono DIE) $\frac{\pi}{3\delta}$ DHq*IH.



Ergo